



Prevalence and demographic characteristics of strabismus in adults

Kourosh Shahraki ^{1,2}, Abolfazl Tahkor ², Fariba Shahraki-Sanavi ², Kianoush Shahraki ^{1,2}, Hasan Okati-Aliabad ², Mahdi Mohammadi ² and Alireza Ansari-Moghaddam ²

¹ Department of Ophthalmology, Zahedan University of Medical Sciences, Zahedan, Iran

² Health Promotion Research center, Zahedan University of Medical Science, Zahedan, Iran

ABSTRACT

Background: Strabismus is a common ocular disorder and a major cause of visual impairment and amblyopia. Its prevalence varies widely across populations and may differ by age, ethnicity, and underlying risk factors. Despite its clinical and psychosocial impact, data on strabismus in Iranian adults remain limited. This study aimed to determine the prevalence of strabismus and describe the demographic distribution of affected adults in southeastern Iran.

Methods: This cross-sectional analysis used baseline data from an adult eye cohort in Zahedan, southeastern Iran. A stratified cluster sampling design, based on socioeconomic status, was used to recruit residents aged 35–70 years between 2015 and 2019. Of 10 016 screened individuals, 9296 met inclusion criteria of Iranian nationality, ≥ 9 months of local residency, and completion of baseline assessments; participants with prior ocular surgery or active ocular disease were excluded. All participants completed a standardized ocular history questionnaire and underwent comprehensive visual assessment, including Snellen visual acuity testing, objective and subjective non-cycloplegic refraction, automated lensometry for habitual correction, external examination, slit-lamp biomicroscopy, and Goldmann applanation tonometry. Ocular alignment was evaluated using cover-uncover and alternate cover tests at distance and near vision. Strabismus was classified based on manifest deviations detected under best correction.

Results: Among 9296 adults, the overall prevalence of strabismus was 2.8% (262 cases). The prevalences of exotropia, esotropia, hypertropia, intermittent exotropia, and intermittent esotropia were 1.4% (129 cases), 0.2% (18 cases), 0.3% (26 cases), 0.8% (77 cases), and 0.1% (12 cases), respectively. Among participants with strabismus, the largest and smallest proportions were in the 45–54-year age group (n = 92, 35.1%) and the 65–70-year age group (n = 14, 5.3%), respectively. Women accounted for 61.8% (n = 162) of all cases, while esotropia cases were equally distributed between women and men. Among affected participants, 34.0% (n = 89) had high school education and 17.6% (n = 46) had university education. The Sistani ethnic group accounted for 53.8% (n = 141) of all cases.

Conclusions: Strabismus affected 2.8% of adults, with exotropia the most common subtype. Among affected participants, the largest proportions were aged 45–54 years, were women, had high school education, and belonged to the Sistani group. Ongoing longitudinal follow-up will clarify incidence, progression, and treatment outcomes, supporting improved clinical decision-making, resource allocation, and long-term visual outcomes.

KEYWORDS

squint, convergent strabismus, divergent strabismus, esodeviation, exodeviation, adults, cohort study, prevalences

Correspondences: Kourosh Shahraki, Department of Ophthalmology, Zahedan University of Medical Sciences, Zahedan, Iran. Email: kourosh.shahyar@gmail.com, ORCID iD: <https://orcid.org/0000-0001-6448-6351>.

How to cite this article: Shahraki K, Tahkor A, Shahraki-Sanavi F, Shahraki K, Okati-Aliabad H, Mohammadi M, Ansari-Moghaddam A. Prevalence and demographic characteristics of strabismus in adults. Med Hypothesis Discov Innov Optom. 2025 Fall; 6(3): 112-120. DOI: <https://doi.org/10.51329/mehdiptometry230>.

Received: 07 March 2025; Accepted: 10 September 2025



Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.



INTRODUCTION

Strabismus is an important ophthalmic disorder, a common ocular condition in childhood, and one of the leading causes of poor vision and amblyopia [1–3]. It is characterized by a deviation of the visual axes in which one eye turns inward, outward, upward, or downward and becomes displaced from the point of fixation [4]. Such deviations may occur in any direction, may be constant or intermittent, and can affect both distance and near vision [4, 5].

Multiple factors may predispose individuals to strabismus, including congenital and hereditary conditions, Mobius syndrome, increased accommodation or convergence leading to esotropia, decreased accommodation or convergence in myopia resulting in exotropia, injury to extraocular nerves or muscles, maternal drug exposure, and other disorders affecting brain function, such as cerebral palsy, Down syndrome, hydrocephalus, brain tumors, and prematurity [4, 6]. Beyond its clinical significance, strabismus is also associated with an elevated risk of psychological, emotional, and social difficulties, which may manifest as depression, anxiety, and impaired social interactions [4, 7, 8].

The prevalence and incidence of strabismus vary worldwide, also within Iran [9–32], with differing rates reported across populations [3, 10, 33–36]. Although esotropia is generally the most common form in most ethnic groups [4, 37], exotropia occurs more frequently than esotropia in certain populations [36, 38], highlighting the need for population- and ethnicity-specific prevalence data.

While such epidemiological investigations are valuable for guiding screening strategies and reducing strabismus-related complications, the condition should also be considered in adults due to its potential association with aging. Studies of this nature facilitate informed clinical decision-making regarding strabismus and other vision disorders that may require surgical or medical intervention. Given the paucity of data on the adult population in Iran, the present study was conducted to determine the prevalence of strabismus and describe the demographic distribution of affected adults in southeastern Iran.

METHODS

The present cross-sectional study utilized baseline data of adults from the PERSIAN Eye Cohort [39] conducted in Zahedan, located in the Sistan and Baluchestan province of southeastern Iran. A total of 10 016 potentially eligible individuals aged 35–70 years residing in the study area were recruited between October 2015 and January 2019. The cohort study received national approval (Grant 700/534) and was endorsed by Zahedan University of Medical Sciences (Grant 93-4-96451). The present investigation was conducted in accordance with the ethical standards outlined in code IR.ZAUMS.REC.1402.107. Written informed consent was obtained from all participants, each of whom was informed of their right to withdraw from the study at any time. Participant data were recorded using unique study identifiers, and strict confidentiality was maintained throughout all stages of the research.

A representative sample of the target population was obtained using a stratified cluster sampling approach, with socioeconomic status serving as the stratifying variable [40, 41]. Initially, several meetings were held with medical experts and public service agencies to categorize areas of Zahedan by socioeconomic status. Next, one district was randomly selected from each of the low, medium, and high socioeconomic strata. All eligible residents within these selected districts were then invited to participate in the study.

Inclusion criteria comprised Iranian nationality, age between 35 and 70 years at the baseline survey, residency in Zahedan for at least nine months, and completion of the baseline survey to enable follow-up assessments. Individuals who did not meet these criteria or who had severe physical or mental illnesses that prevented completion of the questionnaire or attendance at the cohort center [39, 42] were excluded ($n = 9$, 0.1%). Additionally, participants with a history of eye surgery or active ocular disease were excluded ($n = 711$, 7.1%), resulting in a final screened cohort of 9296 subjects.

As part of the eye cohort study, participants completed a self-reported ocular history questionnaire that included information on diabetes, use of spectacles and/or contact lenses, ophthalmologist visits within the past year, dry eye-related symptoms, history of ocular surgery, amblyopia, and family history of glaucoma, retinal detachment, keratoconus, and night blindness. Visual acuity was assessed using a Snellen chart. For participants unable to read letters at a distance of 1 m, a finger-counting test was performed at various shorter distances. If the participant was unable to complete this test, hand movement perception was evaluated at a distance of 30 cm. Participants who still could not perceive hand movements underwent a light perception test, and visual acuity was recorded as “light perception”; if no light perception was detected, it was recorded accordingly.

All participants underwent objective, non-cycloplegic refraction using an autorefractor (NIDEK Auto Ref/Keratometer, Inc., Tokyo, Japan), with the spherical component (in diopters), cylindrical component (in diopter cylinder), and axis of the cylinder (in degrees) recorded for each eye. Additionally, manual objective refraction was performed using a handheld retinoscope (Hein Beta 200 Retinoskop, HEINE Optotechnik, Hersching, Germany) with trial lenses in a trial frame, and any signs of weak red reflex or convergence weakness were documented. For subjective refraction, the trial frame was placed on the participant’s head, with the objective refraction results serving as a starting point. Sphere, cylinder, and axis of cylinder were refined using participants’ responses to the dichromatic and cross-cylinder tests to determine best-corrected distance visual acuity, which was then recorded using individual lenses on the test glasses. For participants wearing habitual near or distance correction, the

optical properties of their spectacles were measured using an automated lensometer (Auto Lensmeter LM-600PD, Nidek, Gamagori, Japan).

A comprehensive external ocular examination was first performed, including inspection of the eyelids and adnexa to detect any eyelid lesions or abnormalities. Pupillary evaluation followed, with specific assessment for pupillary abnormalities and relative afferent pupillary defects [43, 44]. A detailed slit-lamp biomicroscopic examination (SL-D7, Topcon Corporation, Tokyo, Japan) was then conducted for all participants. After instilling topical anesthetic and applying a fluorescein dry strip, intraocular pressure (IOP) was measured in both eyes using a Goldmann applanation tonometer (AT900, Haag-Streit, Koeniz, Switzerland). IOP measurements exceeding 20 mmHg were repeated to ensure reliability. Ocular alignment was subsequently assessed. The cover-uncover and alternate cover tests were performed at distance and near fixation to detect tropias and quantify heterophoria [45]. Testing was performed under best optical correction, with appropriate additional near lenses for presbyopic participants. Each eye was also evaluated individually for the presence of nystagmus. The fixation target consisted of a single Snellen letter (distance or near) presented at one line above the best-corrected visual acuity of the poorer-seeing eye. Participants unable to complete cover testing due to poor cooperation were excluded. For consistency with previous literature, constant tropia, including exotropia, esotropia, hypertropia, and hypotropia, was defined as a unilateral or alternating horizontal or vertical strabismus observed continuously at either distance or near fixation [46, 47]. In the analyses, the unqualified terms “exotropia” and “esotropia” referred to constant deviations and were reported separately from intermittent exotropia and intermittent esotropia.

The data were entered into Microsoft Excel and subsequently imported into IBM SPSS Statistics for Windows, version 23.0 (IBM Corp., Armonk, NY, USA) for analysis. Descriptive statistics were used to characterize the study population and to quantify the distribution of strabismus and its subtypes. Normality of continuous variables was assessed using the Shapiro–Wilk test. Categorical variables, including prevalence and demographic frequencies, were summarized as counts and percentages. Prevalence estimates were calculated as crude proportions using the final analytic sample as the denominator. Continuous variables, such as spherical equivalent and age, were expressed as mean (standard deviation [SD]). All analyses were descriptive, and no inferential hypothesis testing was performed.

RESULTS

A total of 10 016 individuals aged 35–70 years participated in the study. After excluding 720 participants, the final analytic sample comprised 9296 individuals, and the majority were women ($n = 5708$, 61.4%). In the overall population, the mean (SD) spherical equivalent refractive error for the right and left eyes was 0.08 (1.61) diopters and 0.08 (1.53) diopters, respectively.

As summarized in Table 1, the prevalence of exotropia, esotropia, hypertropia, intermittent exotropia, and intermittent esotropia was 1.4%, 0.2%, 0.3%, 0.8%, and 0.1%, respectively. The overall prevalence of strabismus was 2.8%. Among participants diagnosed with strabismus, the highest proportions of exotropia, esotropia, hypertropia, intermittent exotropia, and intermittent esotropia were observed in the 45–54 ($n = 47$, 36.4%), 55–64 ($n = 8$, 44.4%), 35–44 ($n = 10$, 38.5%), 45–54 ($n = 29$, 37.7%), and 55–64 ($n = 5$, 41.7%) years age groups, respectively. Across all strabismus subtypes, the lowest frequencies were consistently observed in the 65–70 years age group. Overall, of the 262 individuals with strabismus, the highest and lowest frequencies were observed in the 45–54 ($n = 92$, 35.1%) and 65–70 ($n = 14$, 5.3%) years age groups, respectively (Table 2).

Among the 262 participants with strabismus, 162 (61.8%) were women and 100 (38.2%) were men; esotropia cases were equally distributed between women and men (Table 2). Regarding educational level, 89 affected participants (34.0%) had high school education and 46 (17.6%) had university education. Among participants with exotropia, hypertropia, intermittent exotropia, and intermittent esotropia, the largest proportions had high school education: 34.1% ($n = 44$), 30.8% ($n = 8$), 32.5% ($n = 25$), and 33.3% ($n = 4$), respectively. Among participants with esotropia, equal proportions were illiterate or had high school education (44.4%, $n = 8$ in each group) (Table 2).

Regarding ethnicity, 141 participants with strabismus (53.8%) were Sistani, 62 (23.7%) were Baloch, and 59 (22.5%) belonged to other ethnic groups. The Sistani group accounted for the largest proportion of cases within each strabismus subtype (Table 2).

Table 1. Prevalence of strabismus subtypes and demographic characteristics of affected participants

Variable	Exotropia ($n = 129$)	Esotropia ($n = 18$)	Hypertropia ($n = 26$)	Intermittent exotropia ($n = 77$)	Intermittent esotropia ($n = 12$)	Total ($n = 262$)
Age (y), Mean $\pm$ SD	52.4 $\pm$ 10.2	54.1 $\pm$ 9.8	51.8 $\pm$ 11.3	53.6 $\pm$ 10.5	55.2 $\pm$ 8.9	-
Sex (Male / Female), n (%)	47 (36.4) / 82 (63.6)	9 (50.0) / 9 (50.0)	8 (30.8) / 18 (69.2)	31 (40.3) / 46 (59.7)	5 (41.7) / 7 (58.3)	100 (38.2) / 162 (61.8)
Ethnicity (Sistani / Baloch / Others), n (%)	67 (51.9) / 32 (24.8) / 30 (23.3)	11 (61.1) / 6 (33.3) / 1 (5.6)	12 (46.2) / 6 (23.0) / 8 (30.8)	44 (57.1) / 14 (18.2) / 19 (24.7)	7 (58.3) / 4 (33.3) / 1 (8.3)	141 (53.8) / 62 (23.7) / 59 (22.5)
Prevalence (%)	1.4	0.2	0.3	0.8	0.1	2.8%

Abbreviations: n , number; y , years; SD, standard deviation; %, percentage.

Table 2. Stratified demographic frequencies of strabismus and its subtypes

Variable	Exotropia, n (%)	Esotropia, n (%)	Hypertropia, n (%)	Intermittent Exotropia, n (%)	Intermittent Esotropia, n (%)	Total, n (%)
Age						
35–44 y	36 (27.9)	2 (11.1)	10 (38.5)	20 (26.0)	2 (16.7)	70 (26.7)
45–54 y	47 (36.4)	7 (38.9)	5 (19.2)	29 (37.7)	4 (33.3)	92 (35.1)
55–64 y	40 (31.0)	8 (44.4)	9 (34.6)	24 (31.2)	5 (41.7)	86 (32.8)
65–70 y	6 (4.7)	1 (5.6)	2 (7.7)	4 (5.2)	1 (8.3)	14 (5.3)
Sex						
Men	47 (36.4)	9 (50.0)	8 (30.8)	31 (40.3)	5 (41.7)	100 (38.2)
Women	82 (63.6)	9 (50.0)	18 (69.2)	46 (59.7)	7 (58.3)	162 (61.8)
Educational level						
Illiterate	34 (26.4)	8 (44.4)	6 (23.1)	22 (28.6)	3 (25.0)	73 (27.9)
Elementary	26 (20.2)	1 (5.6)	7 (26.9)	18 (23.4)	2 (16.7)	54 (20.6)
High School	44 (34.1)	8 (44.4)	8 (30.8)	25 (32.5)	4 (33.3)	89 (34.0)
University	25 (19.4)	1 (5.6)	5 (19.2)	12 (15.6)	3 (25.0)	46 (17.6)
Ethnicity						
Sistani	67 (51.9)	11 (61.1)	12 (46.2)	44 (57.1)	7 (58.3)	141 (53.8)
Baloch	32 (24.8)	6 (33.3)	6 (23.1)	14 (18.2)	4 (33.3)	62 (23.7)
Other	30 (23.3)	1 (5.6)	8 (30.8)	19 (24.7)	1 (8.3)	59 (22.5)
Total	129 (100.0)	18 (100.0)	26 (100.0)	77 (100.0)	12 (100.0)	262 (100.0)

Abbreviations: n, number; %, percentage.

DISCUSSION

In this study, 9296 participants aged 35–70 years were surveyed to determine the prevalence of strabismus. The large population-based sample provides an estimate of the overall prevalence of strabismus in the analytic cohort and describes the demographic distribution of affected participants. The overall prevalence of strabismus was 2.8% (262 cases).

Previous reports show considerable variability depending on age and population characteristics, with estimates ranging from 0.8% in a rural cohort of Indian children [48] to approximately 2%–5% in the general population [4]. The prevalence observed in our study was relatively lower than that reported in most previous studies conducted in Iran [26, 27]. One possible explanation is the age distribution of our participants, as most earlier investigations focused on children and younger individuals [9, 27], and only a few included all age groups [26, 49]. According to current literature, the prevalence of strabismus is 0.80% in Singaporean children aged 6–72 months [13], 2.4% in Hispanic/Latino children [14], 2.5% in African-American children [14], 1.28% in Japanese primary school-aged children 6–12 years (with exotropia and esotropia accounting for 0.69% and 0.28%, respectively) [36], 2.02% in 12.5-year-old Iranian children [27], 1.68% in Iranian seven-year-olds [9], 4.3% in rural Iranian populations [26], 2.8% in Australian six-year-olds [3], 2.3% in English seven-year-olds [50], 4.3% in Canadian children with a mean age of 4.2 years [51], 1.2% in a population-based study of Iranian six-year-olds [52], and 0.81% in a population-based study of children aged 1–12 years in southeastern Brazil [53].

Differences in strabismus prevalence across studies, including ours, likely reflect variation in study design. Many investigations relied on school-based, non-random sampling due to the accessibility of these populations [3, 9, 11, 13, 21, 27, 28, 30, 31, 36, 51, 54–57]. Sample size is another important factor; the present study included a larger sample than most previous studies [26, 27, 29, 58]. Earlier evidence suggests that manifest strabismus tends to appear less frequently in larger population samples [52, 53, 59], although this trend does not apply to heterophoria [60]. It should also be noted that many earlier studies cannot be generalized to the Iranian adult population due to their small sample sizes and narrow age ranges. The present study was designed to address this limitation.

The prevalence of strabismus observed in the present study is comparable to that reported in low-income and lower-middle-income countries, yet lower than estimates from high-income nations [22, 48, 53, 55, 56, 61]. A plausible explanation for the higher prevalence documented in high-income settings is the substantially greater survival rate of low birthweight and premature infants. Advances in neonatal care in these countries raise the survival of premature infants who are at elevated risk for strabismus, thereby contributing to the higher prevalence reported in these populations [62, 63].

Another explanation for the observed variation in strabismus prevalence across studies is the use of differing diagnostic criteria and measurement techniques. Some investigations, including the present one, assessed only manifest strabismus, whereas others included eyes with heterophoria [9, 11, 14, 15, 26, 45, 52, 64, 65]. Diagnostic tests used to quantify ocular deviation likewise vary across studies. Several investigations, similar to ours employed unilateral and alternating cover tests [26, 58, 59], whereas others relied on the Hirschberg test [66] or used a combination of cover testing with corneal light-reflex assessment, with the magnitude of deviation quantified using the Hirschberg method [67].

In a population-based cohort of 3785 Danish adults, the overall prevalence of strabismus was reported at 1.1%, with no significant differences detected by sex or screening status. Within that population, exotropia and esotropia were identified in 0.3% and 0.8% of individuals, respectively, and no vertical deviations were observed [66]. By comparison, in our cohort the prevalence of exotropia, esotropia, and hypertropia was 1.4%, 0.2%, and 0.3%, respectively. Exotropia emerged as the most common horizontal deviation; an observation aligned with several previous reports from Iran. For instance, in a study of 2638

schoolchildren (mean age 12.5 years), exotropia and esotropia were observed in 1.30% and 0.59% of participants, respectively [27]. Similarly, another investigation of 1130 schoolchildren found a strabismus prevalence of 1.9%, with 63.6% of cases classified as exotropic and 36.4% as esotropic [28]. Conversely, some Iranian studies have reported higher overall prevalence rates and differing subtype distributions; one such study documented a strabismus prevalence of 3.1%, with significantly higher rates in girls (4.2%) than in boys (2.0%) [11]. These discrepancies reflect the complex interplay of genetic, environmental, and population-specific factors that shape the epidemiology of strabismus. Consistent with global trends, exotropia tends to predominate in many Asian populations [36, 47, 68, 69] and among individuals of African descent [14], whereas esotropia is more frequently reported in European and Western cohorts [25, 33, 71]. Nonetheless, this pattern is not universal; some studies from Iran [70] and Saudi Arabia [18] have documented a higher prevalence of esotropia, highlighting the regional variability in ocular alignment disorders. Such variation reveals the importance of local epidemiologic data to inform clinical practice, screening strategies, and research priorities. Age is another important factor, as many earlier investigations focused on school-aged children [68, 70], who may not be representative of the general population.

The ratio of exotropia to esotropia in our cohort was 7.2:1. By contrast, a population-based study of Danish adults reported a ratio of 1:2.7, indicating a predominance of esotropia [66]. In pediatric cohorts, ratios of 1.5:1, 9.75:1, and 7:1 have been documented among Asian children [15], Hong Kong children [72], and Singaporean children [13], respectively. The distribution of strabismus types is strongly influenced by the pattern of comorbid refractive error. Exotropia is more commonly associated with myopia, whereas esotropia frequently accompanies hyperopia [46]. Accordingly, the comparatively higher prevalence of exotropia observed in our study may be attributable to the elevated prevalence of myopia within this cohort [42].

A 2016 meta-analysis comprising 23541 participants identified myopia as a significant risk factor for exotropia (odds ratio 5.23), proposing that impaired fusion control at distance arises from delayed accommodation in myopic individuals, thereby reducing accommodative effort [73]. This diminished accommodative drive results in reduced accommodative convergence, and chronic under-engagement of convergence mechanisms is believed to contribute to loss of fusion control, ultimately predisposing individuals to exotropia [74]. Beyond refractive status, several additional independent factors including perinatal exposure to smoking, maternal age, race, and genetic susceptibility, have been implicated in shaping the epidemiology of tropias [71, 75, 76].

In our cohort, women constituted 61.4% of the analytic sample and 61.8% of participants with strabismus. However, evidence from large-scale studies does not support a true sex-based difference in strabismus prevalence. A recent systematic review and meta-analysis by Laughton et al. [71] that synthesized data from 73 population-based and 141 clinic-based studies reported nearly identical prevalence estimates for horizontal strabismus (2.558% in males vs. 2.582% in females), esotropia (1.386% vs. 1.377%), and exotropia (1.035% vs. 1.043%) [71]. Consequently, the higher proportion of women with strabismus in our sample is likely attributable to the overrepresentation of women in the study population. Among participants with esotropia, equal proportions were illiterate or had high school education (44.4%, $n = 8$ in each group). Regarding ethnicity, 53.8% of participants with strabismus were Sistani, 23.7% were Baloch, and 22.5% belonged to other ethnic groups. These apparent differences are likely driven by the unequal distribution of ethnic groups within the study: the Sistani population was disproportionately represented, with 5021 participants contributing to the sample.

This study possesses several important strengths. Foremost, its large, population-based sample provides substantial descriptive information regarding strabismus in the analytic cohort. The use of a standardized, protocol-driven ophthalmic examination minimized inter-examiner variability and reduced the likelihood of measurement error, thereby strengthening the internal validity of the results. An additional advantage of this cohort is its longitudinal design, which will allow future assessments of temporal changes in strabismus prevalence, natural history, and associated risk factors as subsequent waves of data become available from this ongoing study. Nonetheless, several limitations should be acknowledged. The reported prevalence estimates were crude and did not account for the stratified cluster sampling design; therefore, the findings should be interpreted primarily as descriptive estimates for the analytic cohort. In addition, excluding participants with a history of ocular surgery or active ocular disease may have underrepresented previously treated or secondary strabismus and could have resulted in underestimation of prevalence. Some individuals referred for ophthalmic examination also declined participation, introducing possible nonresponse bias. Future research would benefit from the integration of comprehensive ophthalmic records from external clinics, detailed biometric measurements, and extensive environmental and genetic data to further clarify the multifactorial etiology of strabismus. Continued longitudinal follow-up of this cohort is expected to yield valuable insights into incidence patterns, treatment outcomes, and the dynamic interplay between refractive error development and ocular alignment over time.

CONCLUSIONS

This large population-based study provides an updated assessment of strabismus in adults aged 35–70 years. Among 9296 participants, the overall prevalence was 2.8%. Among constant deviations, exotropia was the most common (1.4%), followed by hypertropia (0.3%) and esotropia (0.2%). Intermittent exotropia (0.8%) was more frequent than intermittent esotropia (0.1%). Among participants with strabismus, the largest and smallest proportions were aged 45–54 years and 65–70 years, respectively. Women accounted for 61.8% of cases, reflecting the cohort's demographic structure. Regarding educational and ethnic distributions, 34.0% of affected participants had high school education, and 53.8% belonged to the Sistani group. These findings

describe the demographic distribution of strabismus cases in the analytic cohort. The longitudinal follow-up of this cohort will provide essential evidence on incidence, progression, and treatment outcomes, informing clinical decision-making, referral pathways, and health-system planning. Ultimately, these insights may support more effective resource allocation and improved visual and functional outcomes for adults with strabismus.

ETHICAL DECLARATIONS

Ethical approval: The cohort study received national approval (Grant 700/534) and was endorsed by Zahedan University of Medical Sciences (Grant 93-4-96451). The present investigation was conducted in accordance with the ethical standards outlined in code IR.ZAUMS.REC.1402.107. Written informed consent was obtained from all participants, each of whom was informed of their right to withdraw from the study at any time. Participant data were recorded using unique study identifiers, and strict confidentiality was maintained throughout all stages of the research.

Conflict of interests: None.

FUNDING

The parent PERSIAN Eye Cohort Study received financial support from the Ministry of Health and Medical Education of Iran and the Vice-Chancellor for Research and Technology at Zahedan University of Medical Sciences. No specific funding was received for the present secondary analysis.

ACKNOWLEDGMENTS

We gratefully acknowledge the support of the Ministry of Health and Medical Education of Iran and the Vice-Chancellor for Research and Technology at Zahedan University of Medical Sciences. We also extend our sincere appreciation to the residents of Zahedan for their cooperation and participation, without whom this research would not have been possible.

REFERENCES

1. Suh DW, Shahraki K. Vision Screening Claims for Young Children in the United States. *Pediatrics*. 2023 Sep 1;152(3):e2023062804. doi: 10.1542/peds.2023-062804. PMID: 37605873.
2. Swanson MW, McGwin G. Visual impairment and functional status from the 1995 National Health Interview Survey on Disability. *Ophthalmic Epidemiol*. 2004 Jul;11(3):227-39. doi: 10.1080/09286580490514540. PMID: 15370554.
3. Robaei D, Rose KA, Kifley A, Cosstick M, Ip JM, Mitchell P. Factors associated with childhood strabismus: findings from a population-based study. *Ophthalmology*. 2006 Jul;113(7):1146-53. doi: 10.1016/j.ophtha.2006.02.019. Epub 2006 May 3. PMID: 16675019.
4. Kanukollu VM, Sood G. Strabismus. 2023 Nov 13. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. PMID: 32809617.
5. Shirzadi K, Amirdehi AR, Makateb AL, Shahraki K, Khosravifard K. Studying the effect of tropicamide various concentrations on routine dilation of the pupil. *Biomedical and Pharmacology Journal*. 2015 Dec 28;8(2):885-9. doi: https://dx.doi.org/10.13005/bpj/839.
6. Azonobi IR, Olatunji FO, Adido J, Osayande OO. Vision of strabismic children in Ilorin, Nigeria. *Nigerian Journal of Ophthalmology*. 2008 Nov 12;16(1):12-5. doi: 10.4314/njo.v16i1.11952.
7. Mason A, Joronen K, Lindberg L, Kajander M, Fagerholm N, Rantanen A. Strabismic Adults' Experiences of Psychosocial Influence of Strabismus-A Qualitative Study. *SAGE Open Nurs*. 2024 Sep 5;10:23779608241278456. doi: 10.1177/23779608241278456. PMID: 39246297; PMCID: PMC11378177.
8. Buffenn AN. The impact of strabismus on psychosocial health and quality of life: a systematic review. *Surv Ophthalmol*. 2021 Nov-Dec;66(6):1051-1064. doi: 10.1016/j.survophthal.2021.03.005. Epub 2021 Mar 25. PMID: 33773997.
9. Hashemi H, Yekta A, Jafarzadehpour E, Ostadimoghaddam H, Eshtrati B, Mohazzab-Torabi S, Khabazkhoob M, Soroush S. The prevalence of strabismus in 7-year-old schoolchildren in Iran. *Strabismus*. 2015;23(1):1-7. doi: 10.3109/09273972.2014.999795. Epub 2015 Jan 13. PMID: 25584828.
10. Adejumo OO, Olusanya BA, Ajayi BG. Ocular Disorders among Preschool Children in Southwest Nigeria. *Middle East Afr J Ophthalmol*. 2021 Apr 30;28(1):23-28. doi: 10.4103/meajo.MEAJO_191_19. PMID: 34321819; PMCID: PMC8270023.
11. Faghihi M, Ostadimoghaddam H, Yekta AA. Amblyopia and strabismus in Iranian schoolchildren, Mashhad. *Strabismus*. 2011 Dec;19(4):147-52. doi: 10.3109/09273972.2011.622341. PMID: 22107119.
12. Al-Rowaily MA. Prevalence of refractive errors among pre-school children at King Abdulaziz Medical City, Riyadh, Saudi Arabia. *Saudi J Ophthalmol*. 2010 Apr;24(2):45-8. doi: 10.1016/j.sjopt.2010.01.001. Epub 2010 Apr 2. PMID: 23960874; PMCID: PMC3729549.
13. Chia A, Dirani M, Chan YH, Gazzard G, Au Eong KG, Selvaraj P, Ling Y, Quah BL, Young TL, Mitchell P, Varma R, Wong TY, Saw SM. Prevalence of amblyopia and strabismus in young singaporean chinese children. *Invest Ophthalmol Vis Sci*. 2010 Jul;51(7):3411-7. doi: 10.1167/iovs.09-4461. Epub 2010 Mar 5. PMID: 20207979; PMCID: PMC3979488.
14. Multi-ethnic Pediatric Eye Disease Study Group. Prevalence of amblyopia and strabismus in African American and Hispanic children ages 6 to 72 months the multi-ethnic pediatric eye disease study. *Ophthalmology*. 2008 Jul;115(7):1229-1236.e1. doi: 10.1016/j.ophtha.2007.08.001. Epub 2007 Oct 22. PMID: 17953989; PMCID: PMC4839485.
15. McKean-Cowdin R, Cotter SA, Tarczy-Hornoch K, Wen G, Kim J, Borchert M, Varma R; Multi-Ethnic Pediatric Eye Disease Study Group. Prevalence of amblyopia or strabismus in asian and non-Hispanic white preschool children: multi-ethnic pediatric eye disease study. *Ophthalmology*. 2013 Oct;120(10):2117-24. doi: 10.1016/j.ophtha.2013.03.001. Epub 2013 May 19. PMID: 23697956; PMCID: PMC4848013.

16. Arnold RW. Amblyopia risk factor prevalence. *J Pediatr Ophthalmol Strabismus*. 2013 Jul-Aug;50(4):213-7. doi: 10.3928/01913913-20130326-01. Epub 2013 Apr 2. PMID: 23556991.
17. Han KE, Baek SH, Kim SH, Lim KH; Epidemiologic Survey Committee of the Korean Ophthalmological Society. Prevalence and risk factors of strabismus in children and adolescents in South Korea: Korea National Health and Nutrition Examination Survey, 2008-2011. *PLoS One*. 2018 Feb 14;13(2):e0191857. doi: 10.1371/journal.pone.0191857. PMID: 29444106; PMCID: PMC5812601.
18. Curtis TH, McClatchey M, Wheeler DT. Epidemiology of surgical strabismus in Saudi Arabia. *Ophthalmic Epidemiol*. 2010 Oct;17(5):307-14. doi: 10.3109/09286586.2010.508351. PMID: 20868257.
19. Friedman DS, Repka MX, Katz J, Giordano L, Ibrionke J, Hawse P, Tielsch JM. Prevalence of amblyopia and strabismus in white and African American children aged 6 through 71 months the Baltimore Pediatric Eye Disease Study. *Ophthalmology*. 2009 Nov;116(11):2128-34.e1-2. doi: 10.1016/j.ophtha.2009.04.034. Epub 2009 Sep 16. PMID: 19762084; PMCID: PMC2783780.
20. Garvey KA, Dobson V, Messer DH, Miller JM, Harvey EM. Prevalence of strabismus among preschool, kindergarten, and first-grade Tohono O'odham children. *Optometry*. 2010 Apr;81(4):194-9. doi: 10.1016/j.optm.2009.10.010. PMID: 20346891; PMCID: PMC2847803.
21. Gursoy H, Basmak H, Yaz Y, Colak E. Vision screening in children entering school: Eskisehir, Turkey. *Ophthalmic Epidemiol*. 2013 Aug;20(4):232-8. doi: 10.3109/09286586.2013.808672. PMID: 23865604.
22. Hong JD, Choi EH, Suh S, Bui JH, Storch AM, Walker KR, Shahraiki K, Yanez C, Torres D, Espinoza J, Molina I, Suh DW. UCI EyeMobile Exam Findings from School Children Following on-Site Screening. *Clin Ophthalmol*. 2024 Jun 6;18:1667-1678. doi: 10.2147/OPTH.S460879. PMID: 38860118; PMCID: PMC11164085.
23. Tegegne MM, Fekadu SA, Assem AS. Prevalence of Strabismus and Its Associated Factors Among School-Age Children Living in Bahir Dar City: A Community-Based Cross-Sectional Study. *Clin Optom (Auckl)*. 2021 Apr 8;13:103-112. doi: 10.2147/OPTO.S300124. PMID: 33854396; PMCID: PMC8041646.
24. Wang Y, Zhao A, Zhang X, Huang D, Zhu H, Sun Q, Yu J, Chen J, Zhao X, Li R, Han S, Dong W, Ma F, Chen X, Liu H. Prevalence of strabismus among preschool children in eastern China and comparison at a 5-year interval: a population-based cross-sectional study. *BMJ Open*. 2021 Oct 19;11(10):e055112. doi: 10.1136/bmjopen-2021-055112. PMID: 34667017; PMCID: PMC8527110.
25. Greenberg AE, Mohny BG, Diehl NN, Burke JP. Incidence and types of childhood esotropia: a population-based study. *Ophthalmology*. 2007 Jan;114(1):170-4. doi: 10.1016/j.ophtha.2006.05.072. Epub 2006 Oct 27. PMID: 17070595.
26. Hashemi H, Nabovati P, Yekta A, Ostadimoghaddam H, Behnia B, Khabazkhoob M. The Prevalence of Strabismus, Heterophorias, and Their Associated Factors in Underserved Rural Areas of Iran. *Strabismus*. 2017 Jun;25(2):60-66. doi: 10.1080/09273972.2017.1317820. Epub 2017 May 2. PMID: 28463575.
27. Yekta A, Fotouhi A, Hashemi H, Dehghani C, Ostadimoghaddam H, Heravian J, Derakhshan A, Yekta R, Rezvan F, Behnia M, Khabazkhoob M. The prevalence of anisometropia, amblyopia and strabismus in schoolchildren of Shiraz, Iran. *Strabismus*. 2010 Sep;18(3):104-10. doi: 10.3109/09273972.2010.502957. PMID: 20843187.
28. Yekta A, Hashemi H, Norouzirad R, Ostadimoghaddam H, Nabovati P, Dadbin N, Nirouza F, Shir-Alivand E, Khabazkhoob M. The Prevalence of Amblyopia, Strabismus, and Ptois in Schoolchildren of Dezful. *Eur J Ophthalmol*. 2017 Jan/Feb;27(1):109-112. doi: 10.5301/ejo.5000795. PMID: 27228971.
29. Yekta A, Hashemi H, Ostadimoghaddam H, Haghighi B, Shafiee H, Mehravaran S, Nabovati P, Asharlous A, Khabazkhoob M. Strabismus and Near Point of Convergence and Amblyopia in 4-6 Year-Old Children. *Strabismus*. 2016 Sep;24(3):113-9. doi: 10.1080/09273972.2016.1205103. Epub 2016 Aug 2. PMID: 27485705.
30. Wu J, Wang N. Prevalence and characteristics of amblyopia, strabismus, and refractive errors among patients aged 3-16 years in Shanghai, China: a hospital-based population study. *BMC Ophthalmol*. 2024 Jun 7;24(1):239. doi: 10.1186/s12886-024-03477-8. PMID: 38849757; PMCID: PMC11157824.
31. Agaje BG, Delelegne D, Abera E, Desta K, Girum M, Mossie M, Eshetu D, Hirigo AT. Strabismus prevalence and associated factors among pediatric patients in southern Ethiopia: a cross-sectional study. *J Int Med Res*. 2020 Oct;48(10):300060520964339. doi: 10.1177/0300060520964339. PMID: 33076734; PMCID: PMC7583393.
32. Miyata M, Kido A, Miyake M, Tamura H, Kamei T, Wada S, Ueshima H, Kawai K, Nakao S, Yamamoto A, Suda K, Nakano E, Tagawa M, Tsujikawa A. Prevalence and Incidence of Strabismus by Age Group in Japan: A Nationwide Population-Based Cohort Study. *Am J Ophthalmol*. 2024 Jun;262:222-228. doi: 10.1016/j.ajo.2023.11.022. Epub 2023 Nov 30. PMID: 38040322.
33. Donnelly UM, Stewart NM, Hollinger M. Prevalence and outcomes of childhood visual disorders. *Ophthalmic Epidemiol*. 2005 Aug;12(4):243-50. doi: 10.1080/09286580590967772. PMID: 16033745.
34. Chia A, Roy L, Seenyan L. Comitant horizontal strabismus: an Asian perspective. *Br J Ophthalmol*. 2007 Oct;91(10):1337-40. doi: 10.1136/bjo.2007.116905. Epub 2007 May 2. PMID: 17475715; PMCID: PMC2000992.
35. Pai A, Mitchell P. Prevalence of amblyopia and strabismus. *Ophthalmology*. 2010 Oct;117(10):2043-4; author reply 2044. doi: 10.1016/j.ophtha.2010.06.028. PMID: 20888491.
36. Matsuo T, Matsuo C. The prevalence of strabismus and amblyopia in Japanese elementary school children. *Ophthalmic Epidemiol*. 2005 Feb;12(1):31-6. doi: 10.1080/09286580490907805. PMID: 15848918.
37. Qanat AS, Alsuheili A, Alzahrani AM, Faydhi AA, Albadri A, Alhibshi N. Assessment of Different Types of Strabismus Among Pediatric Patients in a Tertiary Hospital in Jeddah. *Cureus*. 2020 Dec 8;12(12):e11978. doi: 10.7759/cureus.11978. PMID: 33425550; PMCID: PMC7790318.
38. Pan CW, Chen X, Zhu H, Fu Z, Zhong H, Li J, Huang D, Liu H. School-based assessment of amblyopia and strabismus among multiethnic children in rural China. *Sci Rep*. 2017 Oct 17;7(1):13410. doi: 10.1038/s41598-017-13926-8. PMID: 29042639; PMCID: PMC5645399.
39. Alipour F, Hashemi H, Lashay A, Jafari F, Motamed-Gorji N, Jabbarvand Behrouz M, Mirzaei M, Alizade Y, Soleymani MR, Shoja MR, Shahraiki K, Khataminia GR, Poustchi H, Malekzadeh R. PERSIAN Eye Cohort Study (PECS): Design, Methodology. *Arch Iran Med*. 2023 Aug 1;26(8):459-469. doi: 10.34172/aim.2023.70. PMID: 38301109; PMCID: PMC10685732.

40. Tipton E. Stratified sampling using cluster analysis: a sample selection strategy for improved generalizations from experiments. *Eval Rev.* 2013 Apr;37(2):109-39. doi: [10.1177/0193841X13516324](https://doi.org/10.1177/0193841X13516324). Epub 2014 Mar 18. PMID: [24647924](https://pubmed.ncbi.nlm.nih.gov/24647924/).
41. Hennig C, Liao TF. How to find an appropriate clustering for mixed-type variables with application to socio-economic stratification. *Journal of the Royal Statistical Society Series C: Applied Statistics.* 2013 May;62(3):309-69. doi: [10.1111/j.1467-9876.2012.01066.x](https://doi.org/10.1111/j.1467-9876.2012.01066.x).
42. Alipour F, Mohammadzadeh M, Jafari F, Lashay A, Yaseri M, Motamed-Gorji N, Alizadeh Y, Soleimani M, Mirzaei M, Shahraki K, Salimpour S, Shoja MR, Khataminia G, Tahkor A, Tavakoli R, Somi MH, Mansour-Ghanaei F, Joukar F, Ansari-Moghaddam A, Saki N, Hashemi H. Distribution and prevalence of refractive error in Iranian adult population results of the PERSIAN eye cohort study PECS. *Sci Rep.* 2024 Jun 24;14(1):14515. doi: [10.1038/s41598-024-65328-2](https://doi.org/10.1038/s41598-024-65328-2). PMID: [38914645](https://pubmed.ncbi.nlm.nih.gov/38914645/); PMCID: [PMC11196648](https://pubmed.ncbi.nlm.nih.gov/PMC11196648/).
43. Kawasaki AK. Diagnostic approach to pupillary abnormalities. *Continuum (Minneapolis, Minn).* 2014 Aug;20(4 Neuro-ophthalmology):1008-22. doi: [10.1212/01.CON.0000453306.42981.94](https://doi.org/10.1212/01.CON.0000453306.42981.94). PMID: [25099106](https://pubmed.ncbi.nlm.nih.gov/25099106/); PMCID: [PMC10563972](https://pubmed.ncbi.nlm.nih.gov/PMC10563972/).
44. Glazer-Hockstein C, Brucker AJ. The detection of a relative afferent pupillary defect. *Am J Ophthalmol.* 2002 Jul;134(1):142-3. doi: [10.1016/s0002-9394\(02\)01503-9](https://doi.org/10.1016/s0002-9394(02)01503-9). PMID: [12095831](https://pubmed.ncbi.nlm.nih.gov/12095831/).
45. Mestre C, Otero C, Díaz-Doutón F, Gautier J, Pujol J. An automated and objective cover test to measure heterophoria. *PLoS One.* 2018 Nov 1;13(11):e0206674. doi: [10.1371/journal.pone.0206674](https://doi.org/10.1371/journal.pone.0206674). PMID: [30383846](https://pubmed.ncbi.nlm.nih.gov/30383846/); PMCID: [PMC6211723](https://pubmed.ncbi.nlm.nih.gov/PMC6211723/).
46. Sprunger DT, Lambert SR, Hercinovic A, Morse CL, Repka MX, Hutchinson AK, Cruz OA, Wallace DK; American Academy of Ophthalmology Preferred Practice Pattern Pediatric Ophthalmology/Strabismus Panel. Esotropia and Exotropia Preferred Practice Pattern®. *Ophthalmology.* 2023 Mar;130(3):P179-P221. doi: [10.1016/j.ophtha.2022.11.002](https://doi.org/10.1016/j.ophtha.2022.11.002). Epub 2022 Dec 14. PMID: [36526451](https://pubmed.ncbi.nlm.nih.gov/36526451/); PMCID: [PMC10655158](https://pubmed.ncbi.nlm.nih.gov/PMC10655158/).
47. Hashemi H, Pakzad R, Nabovati P, Azad Shahraki F, Ostadimoghaddam H, Aghamirsalim M, Pakbin M, Yekta A, Khoshhal F, Khabazkhoob M. The prevalence of tropia, phoria and their types in a student population in Iran. *Strabismus.* 2020 Mar;28(1):35-41. doi: [10.1080/09273972.2019.1697300](https://doi.org/10.1080/09273972.2019.1697300). Epub 2019 Dec 23. PMID: [31868064](https://pubmed.ncbi.nlm.nih.gov/31868064/).
48. Hegde SA, B S J, V H K. Prevalence of strabismus among school going children in India. *Bioinformation.* 2025 Mar 31;21(3):410-413. doi: [10.6026/973206300210410](https://doi.org/10.6026/973206300210410). PMID: [40599943](https://pubmed.ncbi.nlm.nih.gov/40599943/); PMCID: [PMC12208243](https://pubmed.ncbi.nlm.nih.gov/PMC12208243/).
49. Derakhshan A, Sabermoghaddam A, Abdolalian M, Ganavati SZ, Shakeri MT, Zareei A, Bamdad S. The Strabismus Prevalence in a Random Sample of Iran's Northeastern Population. *Pakistan Journal of Ophthalmology.* 2024;40(1). doi: [10.36351/pjo.v40i1.1688](https://doi.org/10.36351/pjo.v40i1.1688).
50. Williams C, Northstone K, Howard M, Harvey I, Harrad RA, Sparrow JM. Prevalence and risk factors for common vision problems in children: data from the ALSPAC study. *Br J Ophthalmol.* 2008 Jul;92(7):959-64. doi: [10.1136/bjo.2007.134700](https://doi.org/10.1136/bjo.2007.134700). Epub 2008 May 14. PMID: [18480306](https://pubmed.ncbi.nlm.nih.gov/18480306/).
51. Drover JR, Kean PG, Courage ML, Adams RJ. Prevalence of amblyopia and other vision disorders in young Newfoundland and Labrador children. *Can J Ophthalmol.* 2008 Feb;43(1):89-94. doi: [10.3129/i07-187](https://doi.org/10.3129/i07-187). PMID: [18204498](https://pubmed.ncbi.nlm.nih.gov/18204498/).
52. Jamali P, Fotouhi A, Hashemi H, Younesian M, Jafari A. Refractive errors and amblyopia in children entering school: Shahrood, Iran. *Optom Vis Sci.* 2009 Apr;86(4):364-9. doi: [10.1097/OPX.0b013e3181993f42](https://doi.org/10.1097/OPX.0b013e3181993f42). PMID: [19289975](https://pubmed.ncbi.nlm.nih.gov/19289975/).
53. Schaal LF, Schellini SA, Pesci LT, Galindo A, Padovani CR, Corrente JE. The Prevalence of Strabismus and Associated Risk Factors in a Southeastern Region of Brazil. *Semin Ophthalmol.* 2018;33(3):357-360. doi: [10.1080/08820538.2016.1247176](https://doi.org/10.1080/08820538.2016.1247176). Epub 2016 Dec 14. PMID: [27960584](https://pubmed.ncbi.nlm.nih.gov/27960584/).
54. Aring E, Grönlund MA, Andersson S, Hård AL, Ygge J, Hellström A. Strabismus and binocular functions in a sample of Swedish children aged 4-15 years. *Strabismus.* 2005 Jun;13(2):55-61. doi: [10.1080/09273970590922664](https://doi.org/10.1080/09273970590922664). PMID: [16020358](https://pubmed.ncbi.nlm.nih.gov/16020358/).
55. Monaco A, Spadaro A, Sgolastra F, Petrucci A, D'Andrea PD, Gatto R. Prevalence of hyperopia and strabismus in a paediatric population with malocclusions. *Eur J Paediatr Dent.* 2011 Dec;12(4):272-4. PMID: [22185255](https://pubmed.ncbi.nlm.nih.gov/22185255/).
56. Silbert AL, Matta NS, Silbert DI. Incidence of strabismus and amblyopia in preverbal children previously diagnosed with pseudoesotropia. *Am Orthopt J.* 2013;63:103-6. doi: [10.3368/aoj.63.1.103](https://doi.org/10.3368/aoj.63.1.103). PMID: [24141760](https://pubmed.ncbi.nlm.nih.gov/24141760/).
57. Robaei D, Kifley A, Mitchell P. Factors associated with a previous diagnosis of strabismus in a population-based sample of 12-year-old Australian children. *Am J Ophthalmol.* 2006 Dec;142(6):1085-8. doi: [10.1016/j.ajo.2006.06.053](https://doi.org/10.1016/j.ajo.2006.06.053). Epub 2006 Aug 2. PMID: [17157605](https://pubmed.ncbi.nlm.nih.gov/17157605/).
58. Goseki T, Ishikawa H. The prevalence and types of strabismus, and average of stereopsis in Japanese adults. *Jpn J Ophthalmol.* 2017 May;61(3):280-285. doi: [10.1007/s10384-017-0505-1](https://doi.org/10.1007/s10384-017-0505-1). Epub 2017 Feb 22. PMID: [28229342](https://pubmed.ncbi.nlm.nih.gov/28229342/).
59. Yoon KC, Mun GH, Kim SD, Kim SH, Kim CY, Park KH, Park YJ, Baek SH, Song SJ, Shin JP, Yang SW, Yu SY, Lee JS, Lim KH, Park HJ, Pyo EY, Yang JE, Kim YT, Oh KW, Kang SW. Prevalence of eye diseases in South Korea: data from the Korea National Health and Nutrition Examination Survey 2008-2009. *Korean J Ophthalmol.* 2011 Dec;25(6):421-33. doi: [10.3341/kjo.2011.25.6.421](https://doi.org/10.3341/kjo.2011.25.6.421). Epub 2011 Nov 22. PMID: [22131780](https://pubmed.ncbi.nlm.nih.gov/22131780/); PMCID: [PMC3223710](https://pubmed.ncbi.nlm.nih.gov/PMC3223710/).
60. Leone JF, Cornell E, Morgan IG, Mitchell P, Kifley A, Wang JJ, Rose KA. Prevalence of heterophoria and associations with refractive error, heterotropia and ethnicity in Australian school children. *Br J Ophthalmol.* 2010 May;94(5):542-6. doi: [10.1136/bjo.2009.163709](https://doi.org/10.1136/bjo.2009.163709). Epub 2009 Oct 22. PMID: [19850581](https://pubmed.ncbi.nlm.nih.gov/19850581/).
61. Kvamström G, Jakobsson P, Lennerstrand G. Visual screening of Swedish children: an ophthalmological evaluation. *Acta Ophthalmol Scand.* 2001 Jun;79(3):240-4. doi: [10.1034/j.1600-0420.2001.790306.x](https://doi.org/10.1034/j.1600-0420.2001.790306.x). PMID: [11401631](https://pubmed.ncbi.nlm.nih.gov/11401631/).
62. Fieß A, Dautzenberg K, Gißler S, Mildenerberger E, Urschitz MS, Elflein HM, Laspas P, Stoffelns BM, Pfeiffer N, Schuster AK. Prevalence of strabismus and risk factors in adults born preterm with and without retinopathy of prematurity: results from the Gutenberg Prematurity Eye study. *Br J Ophthalmol.* 2024 Oct 22;108(11):1590-1597. doi: [10.1136/bjo-2023-324698](https://doi.org/10.1136/bjo-2023-324698). PMID: [38503479](https://pubmed.ncbi.nlm.nih.gov/38503479/); PMCID: [PMC11503079](https://pubmed.ncbi.nlm.nih.gov/PMC11503079/).
63. Lee DC, Park J, Park HS, Paik HJ, Lee JY, Oh SY, Lee SJ, Lee SY. Characteristic differences between full-term and premature infants with intermittent exotropia. *Sci Rep.* 2024 Sep 19;14(1):21879. doi: [10.1038/s41598-024-72085-9](https://doi.org/10.1038/s41598-024-72085-9). PMID: [39300164](https://pubmed.ncbi.nlm.nih.gov/39300164/); PMCID: [PMC11413369](https://pubmed.ncbi.nlm.nih.gov/PMC11413369/).
64. Su H, Fu J, Chen W, Meng Z, Li L, Dai W, Yao Y. Prevalence of Heterophoria in Tibetan Grade-One Students: The Lhasa Childhood Eye Study. *J Ophthalmol.* 2020 Dec 24;2020:9570908. doi: [10.1155/2020/9570908](https://doi.org/10.1155/2020/9570908). PMID: [33489351](https://pubmed.ncbi.nlm.nih.gov/33489351/); PMCID: [PMC7803105](https://pubmed.ncbi.nlm.nih.gov/PMC7803105/).

65. Sandfeld L, Weihrauch H, Tubaek G, Mortzos P. Ophthalmological data on 4.5- to 7-year-old Danish children. *Acta Ophthalmol*. 2018 Jun;96(4):379-383. doi: [10.1111/aos.13650](https://doi.org/10.1111/aos.13650). Epub 2018 Jan 25. PMID: [29369530](https://pubmed.ncbi.nlm.nih.gov/29369530/).
66. Hultman O, Beth Høeg T, Munch IC, Ellervik C, la Cour M, Andersson Grönlund M, Buch Hesgaard H. The Danish Rural Eye Study: prevalence of strabismus among 3785 Danish adults - a population-based cross-sectional study. *Acta Ophthalmol*. 2019 Dec;97(8):784-792. doi: [10.1111/aos.14112](https://doi.org/10.1111/aos.14112). Epub 2019 Apr 26. PMID: [31025814](https://pubmed.ncbi.nlm.nih.gov/31025814/).
67. Hartness EM, Jiang F, Zamba GKD, Allen C, Bragg TL, Nellis J, Dumitrescu AV, Kardon RH. Automated strabismus evaluation: a critical review and meta-analysis. *Front Neurol*. 2025 Sep 10;16:1620568. doi: [10.3389/fneur.2025.1620568](https://doi.org/10.3389/fneur.2025.1620568). PMID: [41001207](https://pubmed.ncbi.nlm.nih.gov/41001207/); PMCID: [PMC12457155](https://pubmed.ncbi.nlm.nih.gov/PMC12457155/).
68. Matsuo T, Matsuo C, Matsuoka H, Kio K. Detection of strabismus and amblyopia in 1.5- and 3-year-old children by a preschool vision-screening program in Japan. *Acta Med Okayama*. 2007 Feb;61(1):9-16. doi: [10.18926/AMO/32910](https://doi.org/10.18926/AMO/32910). PMID: [17332837](https://pubmed.ncbi.nlm.nih.gov/17332837/).
69. Tanaka A, Ohno-Matsui K, Shimada N, Hayashi K, Shibata Y, Yoshida T, Yamashita M, Tokoro T, Mochizuki M. Prevalence of strabismus in patients with pathologic myopia. *J Med Dent Sci*. 2010 Mar;57(1):75-82. PMID: [20437768](https://pubmed.ncbi.nlm.nih.gov/20437768/).
70. Khorrami-Nejad M, Akbari MR, Khosravi B. The prevalence of strabismus types in strabismic Iranian patients. *Clin Optom (Auckl)*. 2018 Feb 19;10:19-24. doi: [10.2147/OPTO.S147642](https://doi.org/10.2147/OPTO.S147642). PMID: [30214338](https://pubmed.ncbi.nlm.nih.gov/30214338/); PMCID: [PMC6095557](https://pubmed.ncbi.nlm.nih.gov/PMC6095557/).
71. Laughton SC, Hagen MM, Yang W, von Bartheld CS. Gender differences in horizontal strabismus: Systematic review and meta-analysis shows no difference in prevalence, but gender bias towards females in the clinic. *J Glob Health*. 2023 Sep 1;13:04085. doi: [10.7189/jogh.13.04085](https://doi.org/10.7189/jogh.13.04085). PMID: [37651634](https://pubmed.ncbi.nlm.nih.gov/37651634/); PMCID: [PMC10471156](https://pubmed.ncbi.nlm.nih.gov/PMC10471156/).
72. Zhang XJ, Lau YH, Wang YM, Kam KW, Ip P, Yip WW, Ko ST, Young AL, Tham CC, Pang CP, Chen LJ, Yam JC. Prevalence of strabismus and its risk factors among school aged children: The Hong Kong Children Eye Study. *Sci Rep*. 2021 Jul 5;11(1):13820. doi: [10.1038/s41598-021-93131-w](https://doi.org/10.1038/s41598-021-93131-w). PMID: [34226578](https://pubmed.ncbi.nlm.nih.gov/34226578/); PMCID: [PMC8257606](https://pubmed.ncbi.nlm.nih.gov/PMC8257606/).
73. Tang SM, Chan RY, Bin Lin S, Rong SS, Lau HH, Lau WW, Yip WW, Chen LJ, Ko ST, Yam JC. Refractive Errors and Concomitant Strabismus: A Systematic Review and Meta-analysis. *Sci Rep*. 2016 Oct 12;6:35177. doi: [10.1038/srep35177](https://doi.org/10.1038/srep35177). PMID: [27731389](https://pubmed.ncbi.nlm.nih.gov/27731389/); PMCID: [PMC5059633](https://pubmed.ncbi.nlm.nih.gov/PMC5059633/).
74. Kaur K, Gumani B. Exotropia. 2023 Jun 11. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. PMID: [35201713](https://pubmed.ncbi.nlm.nih.gov/35201713/).
75. Yang Y, Wang C, Gan Y, Jiang H, Fu W, Cao S, Lu Z. Maternal smoking during pregnancy and the risk of strabismus in offspring: a meta-analysis. *Acta Ophthalmol*. 2019 Jun;97(4):353-363. doi: [10.1111/aos.13953](https://doi.org/10.1111/aos.13953). Epub 2018 Nov 6. PMID: [30402966](https://pubmed.ncbi.nlm.nih.gov/30402966/).
76. Cotter SA, Varma R, Tarczy-Hornoch K, McKean-Cowdin R, Lin J, Wen G, Wei J, Borchert M, Azen SP, Torres M, Tielsch JM, Friedman DS, Repka MX, Katz J, Ibrionke J, Giordano L; Joint Writing Committee for the Multi-Ethnic Pediatric Eye Disease Study and the Baltimore Pediatric Eye Disease Study Groups. Risk factors associated with childhood strabismus: the multi-ethnic pediatric eye disease and Baltimore pediatric eye disease studies. *Ophthalmology*. 2011 Nov;118(11):2251-61. doi: [10.1016/j.ophtha.2011.06.032](https://doi.org/10.1016/j.ophtha.2011.06.032). Epub 2011 Aug 19. PMID: [21856012](https://pubmed.ncbi.nlm.nih.gov/21856012/); PMCID: [PMC3208120](https://pubmed.ncbi.nlm.nih.gov/PMC3208120/).